

# ANTIMICROBIAL ACTIVITY OF EARTHWORM EXTRACT, *EUDRILUS EUGENIAE* AGAINST FISH BACTERIAL PATHOGENS

**BANSAL NITISH, R. K. GUPTA\* AND SHASHANK**

Department of Zoology,

CCS Haryana Agricultural University, Hisar - 125 004 (Haryana), INDIA

e-mail: rkgupta211066@gmail.com

## INTRODUCTION

Diseases and pollution play an important role in aquaculture. Bacterial disease outbreaks impose a significant constraint on the production of fish and shellfish (Bachere *et al.*, 1995; Verschuere *et al.*, 2000; Gomez *et al.*, 2007). Bacterial pathogens have been reported to cause heavy mortality in both cultured and wild fish species in different parts of the world (Bader *et al.*, 2003; Joseph and Clerk, 2002). The use of anti microbial drugs in agriculture and aquaculture has led to the emergence of antibiotic resistant bacteria (Schwarz *et al.*, 2001; Akinbowale *et al.*, 2006).

Extracting and using biologically active compounds from earthworms has traditionally been practiced by indigenous people throughout the world, more particularly in Asia, including India, Myanmar, China, Korea and Vietnam (Ranganathan, 2006). Earthworms have been known for many centuries as a therapeutic drug source for various diseases in China and other parts of the Far East (Ismail, 2005). Studies on earthworm have shown its antipyretic, antispasmodic, detoxic, diuretic, antihypertensive, antiallergic, anticoagulative, antiasthmatic, spermatocidal, antioxidative, antimicrobial, anticancer, antiulceral and anti-inflammatory activities (Wang *et al.*, 2007; John and Packialakshmi, 2007; Shobha and Kale, 2007; Balamurugan *et al.*, 2009; Cooper and Balamurugan, 2010; Ansari and Sitaram, 2010; Hrzencak *et al.*, 1998).

Recently earthworm protein and its coelomic fluid were reported to exhibit cytolytic, agglutinating, proteolytic, haemolytic, mitogenic, tumorstatic and antibacterial activities (Edwards and Bohlen, 1996; Popovi *et al.*, 2001; Cooper, 2005). Very recently Prakash *et al.* (2007) and Balamurugan *et al.* (2007) have reported that the presence of anti-ulceral and anti-oxidative properties of earthworm paste of *Lampito mauritii*.

A significant number of studies have been carried out by researchers across the globe for assessing and evaluating the antibacterial activity of earthworm powder/paste/extract. However, almost negligible attempts have been made to study this extra-ordinary property of earthworms in fish health management. In order to scientifically analyze some of the ethnomedical uses of earthworm, the present investigation was undertaken to analyze the antibacterial properties of earthworm extract and its use in fish health management.

## MATERIALS AND METHODS

The study was carried out at Fish Biotechnology Laboratory and Aquarium Room, Department of Zoology, Chaudhary Charan Singh Haryana Agricultural University, Hisar. The following materials and methods were used during the research work.

### Isolation and identification of bacterial strains from fish

#### Collection of material

## ABSTRACT

The present study was conducted to examine antimicrobial activity of earthworm extract, *Eudrilus eugeniae* against fish bacterial pathogens. 8 bacterial strains (6 gram negative and 2 gram positive) were isolated and identified from diseased fish. The antimicrobial activity of extract from *Eudrilus eugeniae* was determined under *in vivo* study by using antimicrobial well diffusion assay. The earthworm extract showed antibacterial activity against isolated pathogenic bacterial strains. The maximum zone of inhibition ( $21.00 \pm 0.57$  mm) was shown against *Aeromonas hydrophila* by earthworm extract. The study establishes that earthworm extract from *Eudrilus eugeniae* has great antibacterial potential and can be used to suppress bacterial infection in fishes.

## KEY WORDS

Bacteria  
Common carp  
*Eudrilus eugeniae*, zone of inhibition  
Antimicrobial activity

**Received : 23.01.2015**

**Revised : 22.01.2016**

**Accepted : 04.02.2016**

**\*Corresponding author**

Healthy and infected fish (*Cyprinus carpio*) were collected from fish farms/ponds in Hisar (Haryana) and brought to the Fish Biotechnology Laboratory of the department. The external symptoms of the infected fish were recorded. The specimens were kept under laboratory conditions for isolation and the identification of bacterial strains. The earthworms, *Eudrilus eugineae* were collected from Vermicomposting Unit, Department of Zoology, CCSHAU, Hisar.

#### Culture of bacteria

The tissues and organs (skin lesions, gills and intestine) were dissected out and homogenized by the homogenizer in the glass tube. The supernatant was spread over the nutrient agar (NA) medium under aseptic conditions. The plates were incubated in B.O.D at  $30 \pm 2^\circ\text{C}$  for 18-24 hours. Growth on NA plate was observed after 18-24 hours. Pure colonies of bacteria were obtained by further sub-culturing single colonies on NA plates. (Gerhardt *et al.*, 1994)

#### Identification of bacteria

Isolated pure culture bacteria were subjected to a number of important biochemical tests (primary and secondary tests for identification described by Krieg and Holt (1984) in "Bergey's Manual of Determinative Bacteriology". The identification of bacteria was done with the help of Computer program, PIBWin (Website: [www.soton.ac.uk/tnb/pib.htm](http://www.soton.ac.uk/tnb/pib.htm)).

#### In vivo pathogenicity test

Healthy fish (common carp) weighing about 20-25g were procured from the fish farms and brought to the laboratory. The fish were acclimated for one week in the laboratory. The fish were fed a normal recommended commercial diet. *In vivo* pathogenicity test was carried out on healthy fish by method as described by Keskin *et al.* (2004).

Pure culture of the isolated and identified bacteria with known viable count were injected intraperitoneally (i.p.) into fish @ 0.5 ml per fish. The control fish were inoculated with normal saline. The experiment was done in triplicates with 10 fish in each tub in each replicate. The symptoms of disease appearance were examined and incubation period of different bacteria in fish and longevity of fish were recorded.

#### Preparation of earthworm paste (Balamurugan *et al.* 2007)

Earthworms (*Eudrilus eugineae*) were obtained from Vermicomposting Unit of the Department of Zoology. Sexually mature clitellated worms were washed with running tap water and then fed with wet blotting paper for 18-20 hours for gut clearance. The gut cleared worms were again washed with distilled water. The worms were kept in plastic trough covered tightly with polythene cover and exposed to sunlight for three days to kill the earthworms. Mucus and coelomic fluid that oozed out formed a brown coloured paste called earthworm paste. After mastication of whole earthworm obtained with mucus and coelomic fluid, a brown coloured paste was formed.

#### Well diffusion technique (Gram and Melchiorson, 1996)

Nutrient agar (NA) was cooled at  $45^\circ\text{C}$  and inoculated with the target organism to a final density of  $10^7$  cells /mL agar media and was poured into the Petri dishes as described by Gram and Melchiorson, (1996). The plates were kept in Laminar Air Flow for 30 min. to allow solidification. With the

help of cork-borer, three wells were prepared in each solidified agar plate and were sealed with water agar. The earthworm extract (50 ml) was inoculated in wells ( $r = 0.25$  cm) with the help of a micropipette. The plates were then incubated at  $32 \pm 1^\circ\text{C}$  in B.O.D. incubator and observations were recorded for zone of inhibition around the wells after 24h. The clear zone of inhibition was measured using a centimeter scale and the observations were recorded.

#### Statistical analysis

The obtained results were analyzed statistically using completely randomized design (CRD) following Snedecor and Cochran (1989).

## RESULTS

Isolation of bacterial strains was done from healthy and diseased *Cyprinus carpio* collected from fish farms/ ponds in Hisar. The diseased fish showed symptoms like hemorrhages on head and lateral side of the body, skin erosions, Fin and tail erosion, ulcers and rotting of gill lamella. Isolated pure cultures of bacteria were subjected to a number of important biochemical tests. Eight bacterial strains were isolated and identified from *Cyprinus carpio* (Table 1).

#### Longevity of *Cyprinus carpio* inoculated with different pathogenic bacteria

Table 2 shows the longevity in days of fish infected with different

**Table 1: Incubation period of different pathogenic bacteria for the appearance of disease symptoms in *Cyprinus carpio***

Pathogenic bacteria	Incubation period (in days) <sup>a</sup> of bacteria for the appearance of disease symptoms
Gram negative	
<i>Aeromonas hydrophila</i>	3.34 $\pm$ 0.33
<i>Escherichia coli</i>	7.64 $\pm$ 0.57
<i>Enterobacter aerogens</i>	8.32 $\pm$ 0.33
<i>Shigella spp.</i>	8.47 $\pm$ 0.33
<i>Pseudomonas fluorescens</i>	6.75 $\pm$ 0.68
<i>Pseudomonas aeruginosa</i>	8.13 $\pm$ 0.33
Gram positive	
<i>Micrococcus luteus</i>	7.00 $\pm$ 0.33
<i>Staphylococcus aureus</i>	5.07 $\pm$ 0.57
CD Value (P < 0.05)	0.24

a = Mean  $\pm$  S.E.; N = 30 (10 fish x 3 replications)

**Table 2: Longevity of *Cyprinus carpio* injected with different pathogenic bacteria**

Pathogenic bacteria	Longevity (in days) <sup>a</sup>
Gram negative	
<i>Aeromonas hydrophila</i>	8.66 $\pm$ 0.33
<i>Escherichia coli</i>	12.14 $\pm$ 0.57
<i>Enterobacter aerogens</i>	15.52 $\pm$ 0.33
<i>Shigella spp.</i>	16.37 $\pm$ 0.33
<i>Pseudomonas fluorescens</i>	11.53 $\pm$ 1.20
<i>Pseudomonas aeruginosa</i>	13.70 $\pm$ 0.57
Gram positive	
<i>Micrococcus luteus</i>	13.25 $\pm$ 0.33
<i>Staphylococcus aureus</i>	10.38 $\pm$ 0.57
CD (P < 0.05)	0.28

a = Mean  $\pm$  S.E.; N = 30 (10 fish x 3 replications)

pathogenic bacteria. The fish injected with *A. hydrophila* survived for about 8 days post inoculation. The *A. hydrophila* was the most pathogenic bacterium among the isolated strains. *P. fluorescens* and *S. aureus* was also quite pathogenic as fish could live just for about 10-11 days post inoculation and mortality observed was 80.25% (Table 2). Among pathogenic bacteria maximum longevity was seen in *Shigella* spp. wherein fish survived for about 16 days. This was followed by fish injected with *E. aerogens* where longevity observed was about 15 days.

#### Antibacterial activity shown by earthworm species against isolated pathogenic bacteria.

##### *In vitro* antagonism test

The earthworm species i.e. *Eudrilus eugineae*, was tested for their antagonistic behavior towards pathogenic bacterial strains using well diffusion technique. The results of the *in vitro* antagonism tests are presented in Table 3.

The maximum values of zone of inhibition against the fish bacterial pathogens were given by *Eudrilus eugineae* in the composition of 1:1(Extract: Ethanol)followed by pure extract and 1:2 (Extract: Ethanol).The mean values of the zone of inhibition observed for *Eudrilus eugeneia* against isolated pathogenic bacteria are shown in Table 3 (Fig. 1-8).

The maximum zone of inhibition observed for *Eudrilus eugeneia* was 21.00 mm against *A. hydrophila* and the minimum was 14.00 mm against *Shigell* sp. in gram negative bacteria. In case of gram positive bacteria the maximum value observed was 19.66 mm against *S. aureus*.

## DISCUSSION

The present investigation began with screening of antibacterial activity of different earthworm species against isolated bacterial strains from healthy and diseased fish. 6 gram negative and 2 gram positive bacterial strains were isolated from fish, *Cyprinus carpio*. The number of bacterial isolates can be correlated with studies of various scientists who reported that intestinal micro flora of aquatic animals consists mainly of Gram-negative aerobic, obligate anaerobic and facultative anaerobic bacteria, the composition of which may change with environmental stresses (Ringo and Strøm, 1994; Ringo et al., 1997; Kennedy et al., 1998), diet (Munro et al., 1993; Douillet and Langdon, 1994; Gildberg et al., 1995;) and fish age (Bergh

et al., 1994; Prayitno and Latchford, 1995; Olafsen, 2001). The isolated bacterial strains belonged to *Aeromonas* spp, *Pseudomonas* spp. and members of the family Enterobacteriaceae (Anand et al., 2011; Hassan et al., 2011; Satish et al., 2013). These bacteria

dominate micro flora of freshwater species (Ugajin, 1979; Sugita et al., 1988; Sakata, 1990; Sugita et al., 1991; Ringo et al., 1995). The motile aeromonas group, especially *A. hydrophila*, affects a wide variety of freshwater fish species and occasionally marine fish (Larsen and Jensen, 1977). *A. hydrophila* was found to be most pathogenic to fish as it took the least time to show the disease symptoms in fish. Harikrishnan et al. (2003) observed the pathogenicity induced by *A. hydrophila* when injected @10<sup>8</sup>cfu/ml in common carp, *Cyprinus carpio* and found hemorrhagic spot at the site of injection on the 7th day, whereas in the present study the lesions appeared on the 4<sup>th</sup> day of the injection in the fish *Cyprinus carpio*. The prophylactic and therapeutic control of the bacterial diseases is based on oral administration of antibiotics. However, such treatment may cause the development of resistant bacteria (Aoki et al., 1985), yield residues in fish and introduce potential hazard to public health and to the environment. A new approach method, that is gaining acceptance within the industry, is the use of earthworm extract to control potential pathogens.

In the present study *in vitro* antagonism test following well diffusion assay was conducted to determine the zone of inhibition of earthworm extract against various fish pathogens. The maximum values of zone of inhibition against the fish bacterial pathogens were given by earthworm extract: ethanol (1:1) followed by pure extract and earthworm extract: ethanol (1:2) among all the composition used. The results of the study can be associated to study of Vasanthi et al. (2013) who worked onantimicrobial activity of earthworm paste *Eudrilus eugeniae* and found that earthworm paste at a dose of 100µL was able to inhibit the growth of bacteria of *S. aureus* at a maximum level compared to other bacteria; the growth of fungal *Candida albicans* was much inhibited. Kathireswari et al. (2014) observed the similar results and revealed that coelomic fluid of earthworms *Lampito mauritii* and *Megascolex konkanensis* have high antimicrobial activity against *A. hydrophila*, *Bacillus subtilus*, *Vibrio para haemolyticus*.

Similar results were observed by Mathur et al. (2010) who



Figure 1: Zone of inhibition against *A. hydrophila*.



Figure 2: Zone of inhibition against *E. aerogens*

**Table 3: Inhibition zones (in mm) of *Eudrilus eugineae* against different pathogenic bacteria.**

Bacterial strains	Pure Extract	Extract:Ethanol1:1	Extract:Ethanol 1:2
<i>Aeromonas hydrophila</i>	20.33 ± 0.33	21.00 ± 0.57	18.66 ± 0.33
<i>Staphylococcus aureus</i>	17.66 ± 0.33	19.66 ± 0.33	14.66 ± 0.66
<i>Pseudomonas aeruginosa</i>	17.33 ± 0.33	17.33 ± 0.33	15.33 ± 0.33
<i>Pseudomonas florescens</i>	18.33 ± 0.33	19.33 ± 0.33	15.00 ± 0.57
<i>Micrococcus luteus</i>	13.00 ± 0.57	15.00 ± 0.57	12.66 ± 0.33
<i>Escherichia coli</i>	19.00 ± 0.57	20.66 ± 0.33	15.33 ± 0.33
<i>Enterobacteraerogens</i>	17.66 ± 0.33	18.33 ± 0.33	14.66 ± 0.33
<i>Shigella spp.</i>	11.33 ± 0.88	14.00 ± 0.57	9.00 ± 0.57
C.D. (Pd <sup>0.05</sup> )	1.51	1.33	1.38

a = Mean ± S.E.

**Figure 3: Zone of inhibition against *E. coli*****Figure 4: Zone of inhibition against *P. fluorescens*****Figure 5: Zone of inhibition against *S. aureus*****Figure 6: Zone of inhibition against *P. aeruginosa***

studied antimicrobial activity of earthworm extracts of various solvents and found that 95% ethanol extract of earthworm was potent antibacterial agent against *Streptococcus pyogenes* and antifungal agent against *Candida albicans*. Istiqumah *et al.* (2012) studied to determine the inhibitory of earthworm (*Lumbricus rubellus*) extract (ECT) and encapsulated earthworm extract (ECT-t) as poultry feed additive against some pathogenic bacteria and found that *in vitro* antibacterial activity was performed using dilution method against *Escherichia coli*, *Staphylococcus aureus*, *Salmonella pullorum* and *Pseudomonas aeruginosa*. Elif Ozlem and Cotuk

(2008) worked on antibacterial and haemolytic activity of the coelomic fluid of *Dendrobaena aveneta* (Oligochaeta, Lumbricidae) living in different localities and concluded that coelomic fluid of Beykoz earthworms was a potential agent, which can be used as an alternative drug, since this coelomic fluid was effective against bacteria. The present study indicates that the earthworm extract has good antibacterial potency. Similar results were reported by Shobha and Kale (2008) who found that coelomic fluid has antibacterial activity against bacteria. Bhorgin and Uma (2014) also found that 95% ethanol extract of earthworm was potent antimicrobial agent against



Figure 7: Zone of inhibition against *M. luteus*



Figure 8: Zone of inhibition against *Shigella spp.*

*A. hydrophila* and antifungal agent against *C. albicans*. Based on the living habitats of earthworms, it is rational to think that there are some bioactive compounds that showed effective antimicrobial agents in earthworm's skin. Cho *et al.* (1998) identified the first antimicrobial peptide (lumbricin I) from the earthworm, *Lumbricus rubellus*. Lumbricin I is considered as a proline-rich antimicrobial peptide containing 62 amino acids including proline (15%) with a molecular weight of 7231 Da. Lumbricin I showed antimicrobial activity *in vitro* against a broad spectrum of microorganisms without hemolytic activity. Engelmann *et al.* (2004) and Balamurugan *et al.* (2008) found that earthworm coelomic fluid contains biologically active molecules and leukocytes that participate in phagocytosis, encapsulation and killing of HeLa, HEP-2, PC-12 and PA317 cells *in vitro*.

In conclusion, it can be stated that earthworm extract can reduce the incidence and duration of diseases. The application of earthworm extract in aquaculture shows promise but needs considerable efforts of research.

## ACKNOWLEDGMENT

The authors are thankful to Professor and Head, Department of Zoology, College of Basic Sciences and Humanities, CCSHAU, Hisar for providing facilities and support rendered. The authors are also thankful to Department of Science and Technology, Min. of Science and Technology for providing financial support to carry out this research.

## REFERENCES

- Akinbowale, O. L., Peng, H. and Barton, M. D. 2006. Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. *J. App. Microbiol.* **100**: 1103-1113.
- Anand, J., Thomas, A. Akhila and Salini, M. P. 2011. Fluctuations in the microbial load at certain nonspecific immune sites of macrobrachium rosenbergii supplemented with oicumum sanctum. *The Ecoscan.* **1**: 389-392.
- Ansari, A. A. and Sitaram, K. 2010. An investigation on the anti-microbial and anti-fungal properties of earthworm powder obtained from *Eiseniafoetida*. *Int. J. microbio. res.* **1(2)**: 62-66.
- Aoki, T., Kanazawa, T. and Kitao, T. 1985. Epidemiological surveillance of drug-resistant *Vibrio anguillarum* strains. *Fish Pathol.*

**20**: 199-208.

Bachere, E., Mialhe, E., Noël, D., Boul, V., Morvan, A. and Rodrigues, J. 1995. Knowledge and research prospects in marine mollusc and crustacean immunology. *Aquacult.* **132**: 17-32.

Bader, J. A., Shoemaker, C. A. and Klesius, P. H. 2003. Rapid detection of columnaris disease in channel catfish (*Ictalurus punctatus*) with a new species specific 16S rRNA gene based PCR primer for *Flavobacterium columnare*. *J. Microbiol. Met.* **52(2)**: 209-219.

Balamurugan, M., Parthasarathi, K., Cooper, E. L. and Ranganathan, L. S. 2007. Earthworm paste (*Lampitomaauritii*, Kinberg) alters inflammatory, oxidative, haematological and serum biochemical indices of inflamed rat, *European Review for Medical and Pharmacological Sciences.* **11**: 77-90.

Balamurugan, M., Parthasarathi, K., Cooper, E. L. and Ranganathan, L. S. 2009. Anti-inflammatory and anti-pyretic activities of earthworm extract- *Lampitomaauritii* (Kinberg). *J. Ethnopharm.* **121(2)**: 330-332.

Balamurugan, M., Parthasarathi, K., Cooper, E. L. and Ranganathan, L. S. 2008. Hypothetical mode of action of earthworm extract with hepatoprotective and antioxidant properties. *J. Zhejiang Univ. Sci. B.* **9(2)**: 141-147.

Bergh, O., Naas, K. E. and Harboe, T. 1994. Shift in the intestinal microflora of Atlantic halibut (*Hippoglossus hippoglossus*) larvae during first feeding. *Can. J. Fish. Aquat. Sci.* **51**: 1899-1903.

Bhorgin, A. J. and Uma, K. 2014. antimicrobial activity of earthworm powder (*Lampito mauritii*). *Int. J. Curr. Microbiol. App. Sci.* **3(1)**: 437-443.

Cho, J. H., Park, C. B., Yoon, Y. G. and Kim, S. C. 1998. Lumbricin I, a novel proline-rich antimicrobial peptide from the earthworm: purification, cDNA cloning and molecular characterization. *Biochim. Biophys. Acta.* **1408**: 67-76.

Cooper, E. L. 2005. CAM, eCAM, bioprospecting: the 21st century pyramid, *Evidence Based Complementary Ancient Medicines.* **2**:125-127.

Cooper, E. L. and Balamurugan, M. 2010. Unearthing a source of medicinal molecules. *J. Ethnopharm.* **10**: 70-76.

Douillet, P. A. and Langdon, C. J. 1994. Use of a probiotic for the culture of larvae of the Pacific oyster (*Crassostrea gigas* Thunberg). *Aquacult.* **119**: 25-40.

Edwards, C. A. and Bohlen, P. J. 1996. Biology and Ecology of Earthworms (third edition), Chapman and Hall, London.

ElifOzlem Arslan, A. and Aypýn, C. 2008. Antibacterial and Hemolytic Activity of the Coelomic Fluid of *Dendrobaenaveneta*(Oligochaeta, Lumbricidae) Living in Different Localities. *IUFS J Biol* 2008. **67(1)**: 23-32.

Engelmann, P., Kiss, J., Cooper, E. L. and Nemeth, P. 2004. Earthworm

- leukocytes kill HeLa, HEp-2, PC-12 and PA317 cells in vitro. *J. Biochem. Biophys. Methods*. **61**: 215-228.
- Gerhardt, P., Murray, R. G. E., Wood, W. A. and Kreig, N. R. 1994.** methods for general and molecular bacteriology. *American society for microbiology press, Washington, DC.*
- Gildberg, A., Johansen, A. and Bogwald, J. 1995.** Growth and survival of Atlantic salmon (*Salmo salar*) fry given diets supplemented with fish protein hydrolysate and lactic acid bacteria during a challenge trial with *Aeromonas salmonicida*. *Aquacult.* **138**: 23-34.
- Gómez, R., Balcázar, J. L. and Shen, M. A. 2007.** Probiotics as Control Agents in Aquaculture. *J. Ocean Uni. China*. **6(1)**: 76-79.
- Gram, L. and Melchiorson, J. 1996.** Interaction between fish spoilage bacteria *Pseudomonas* sp. and *Shewanellaputrefaciens* in fish extracts and on fish tissue. *J. Appl. Bacteriol.* **80**: 589-595.
- Harikrishan, R., Nishrani, M. and Balasundaram, C. 2003.** Hematological and Biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophila* infection. *Aquacult.* **221(1-4)**: 41-50.
- Hassan, S., Qureshi, T. A., Ahmad, B. and Manohar, S. 2011.** Microflora of ice in fish carrying crates from certain fish markets of bhopal. *The Ecoscan*. **5(1&2)**: 81-83.
- Hrzenjak, T., Popovic, M., Bozic, T., Grdisa, M., Kobrehel, D. and Rudman, L. 1998.** Fibrinolytic and anticoagulative activities from the earthworm *Eiseniafoetida*. *Comp. Biochem. and Physiol.* **119**: 825-832.
- Ismail, S. A. 2005.** The Earthworm Book. Other India Press, Apusa, Goa, p. 101.
- Istiqomah, L., Herdian, H., Damayanti, E., Hayati, S. N. and Julendra, H. 2012.** Inhibitory of Encapsulated Earthworm Extract (*Lumbricusrubellus*) on Pathogenic Bacteria *In Vitro*. Media Peternakan, April. pp. 1-8.
- John, S. A. and Packialakshmi, N. 2007.** Studies on antimicrobial activity of *Perionyxeccavatus*. *Pakistan J. biological sciences*. **10(19)**: 3482-3484.
- Joseph, A. and Clerk, G. C. 2002.** Infestation of fish-culturing communities with fish-borne bacteria: The Ghanain case. *Int. J. Environ. Health Res.* **12(3)**: 277-282.
- Kathireswari, P., Alakesan, A., Abirami, P. and Sangeetha, P. 2014.** Antimicrobial activity of earthworm coelomic fluid against diseases causing microorganisms. *Int. J. Curr. Microbiol. App. Sci.* **3(8)**: 608-613.
- Kennedy, S. B., Tucker, J. W., Thoresen, M. and Sennett, D. G. 1998.** Current methodology for the use of probiotic bacteria in the culture of marine fish larvae. *Aquaculture 98, World Aqua. Soc., Baton Rouge*. p. 286.
- Keskin, O., Secer, S., Izgor, M., Turkyilmaz, S. and Makaosya, R. S. 2004.** *Edwardsiella ictaluri* Infection in Rainbow Trout (*Oncorhynchus mykiss*). *J. Vet. Anim. Sci.* **28**: 649-653.
- Krieg, N. R. and Holt, J. G. 1984.** Identification of bacteria In: *Bergey's manual of systematic bacteriology. Vol. I Williams and Wilkins Baltimore, London*. pp. 24-26.
- Larsen, J. L. and Jensen, N. J. 1977.** An *Aeromonas* species implicated in ulcer-disease of the cod (*Gadusmorhua*). *Nord Vet. Med.* **29**: 199-211.
- Mathur, A., Satish, K., Verma, Rakshanda Bhat, Santosh, K. Singh, Archana, P., Prasad, G. B. K. S. and Dua, V. K. 2010.** Antimicrobial Activity of Earthworm Extracts. *Intjou of Pharma Scie.* **2(4)**: 364-370.
- Munro, P. D., Birkbeck, T. H. and Barbour, A. 1993.** Influence of rate of bacterial colonisation of the gut of turbot larvae on larval survival. In *Fish Farming Technology - Proceedings of the First International Conference of on Fish Farming Technology*. Reinerstein, H., Dahle, L. A., Jørgensen, L., and Tvinnereim, K. (eds.). A.A. Balkema, Rotterdam pp. 85-92.
- Olafsen, J. A. 2001.** Interactions between fish larvae and bacteria in marine aquaculture. *Aquacult.* **200**: 223-247.
- Popovi, M., Mihaela, T. H. R., Babi, T., Kos, J. and Grdisa, M. A. 2001.** Effect of earthworm (G-90) extract on formation and lysis of clots originated from venous blood of dogs with cardiopathies and with malignant tumors, *Pathology Oncology Research*. **7**: 197-202.
- Prakash, M., Balamurugan, M., Parthasarathi, K., Gunasekaran, G., Cooper E. L. and Ranganathan, L. S. 2007.** Anti-ulceral and anti-oxidative properties of "earthworm paste" of *Lampitomaureitii* (Kinberg) on *Rattusnorvegicus*, *European Review for Medical and Pharmacological Sciences*. **11**: 9-15.
- Prayitno, S. B. and Latchford, J. W. 1995.** Experimental infections of crustaceans with luminous bacteria related to *Photobacterium* and *Vibrio*. *Aquacult.* **132**: 105-112.
- Ranganathan, L. S. 2006.** Vermibiotechnology—From Soil Health to Human Health. *Agrobios, Jodhpur, India*.
- Ringo, E. and Strøm, E. 1994.** Microflora of Arctic charr, *Salvelinus alpinus* (L.): gastrointestinal microflora of free-living fish and effect of diet and salinity on intestinal microflora. *Aquacult Fish Manage.* **25**: 623-629.
- Ringo, E., Olsen, R. E., Overli, O. and Lovik, F. 1997.** Effect of dominance hierarchy formation on aerobic microbiota associated with epithelial mucosa of subordinate and dominant individuals of Arctic charr, *Salvelinus alpinus* (L.). *Aquac. Res.* **28**: 901-904.
- Ringo, E., Strøm, E. and Tabachek, J. A. 1995.** Intestinal microflora of salmonids: a review. *Aquac. Res.* **26**: 773-789.
- Sakata, T. 1990.** Microflora in the digestive tract of fish and shellfish. In *Microbiology of Poecilotherms*. Lesel, R. (ed.). Elsevier, Amsterdam pp. 171-176.
- Satish, P. V. V., Sree Rekha, N., Sunita, K., Somaiah, K. and Kanaka Raju, A. 2013.** Prevalence of fin and tail rot disease in nursery ponds of guntur district, andhra pradesh, India. *The Ecoscan*. **7(3&4)**: 95-99
- Schwarz, S., Kehrenberg, C. and Walsh, T. R. 2001.** Use of antimicrobial agents in veterinary medicine and food animal production. *Intern. J. Antimicrob. Agent.* **17**: 431-437.
- Shobha, S. V. and Kale, R. D. 2008.** *In vitro* studies on control of soil-borne plant pathogens by earthworm *Eudrilus eugeniae* exudates. Green Pages. [http://www.ecoweb.com/ editorial/ 080106.html](http://www.ecoweb.com/editorial/080106.html).
- Shobha, S. V. and Kale, R. 2007.** Antimicrobial potency of *Eudrilluseugeniae* on certain plant pathogen//abstract 8<sup>th</sup> Int. Symp.on earthworm ecology, Krakow, Poland, 4-9 sept, 2006, p. 245.
- Snedecor, G. and Cochran, W. 1989.** *Statistical Methods* (8<sup>th</sup> edition). The Iowa State University Press.
- Sugita, H., Miyajima, C. and Deguchi, Y. 1991.** The vitamin B-12 producing ability of the intestinal microflora of freshwater fish. *Aquacult.* **92**: 267-276.
- Sugita, H., Tsunohara, M., Ohkoshi, T. and Deguchi, Y. 1988.** The establishment of intestinal microflora in developing goldfish (*Carassius auratus*) of culture ponds. *Microbial Ecol.* **15**: 333-344.
- Ugajin, M. 1979.** Studies on the taxonomy of major microflora on the intestinal contents of salmonids. *B. Jpn. Soc. Sci. Fish.* **45**: 721-731.
- Vasanthi, K., Chairman, K. and Ranjit Singh, A. J. A. 2013.** Antimicrobial activity of earthworm paste *Eudrilluseugeniae*. *African J. Environmental Science and Technology*. **7(8)**: 789-793.
- Verschuere, L., Rombaut, G., Sorgeloos, P. and Verstraete, W. 2000.** Probiotic bacteria as biological control agents in aquaculture. *Microbiol. Molec. Biol. Rev.* **64**: 655-671
- Wang, C., Sun, Z., Liu, Y., Zhang, X. and Xu, G. 2007.** A novel antimicrobial vermipeptide family from earthworm *Eiseniafetida*. *European J. Soil Biology*. **43**: 127-134.